

# Quantifiers for quantum logic

Chris Heunen

April 23, 2009

## Abstract

We consider categorical logic on the category of Hilbert spaces. More generally, in fact, any pre-Hilbert category suffices. We characterise closed subobjects, and prove that they form orthomodular lattices. This shows that quantum logic is just an incarnation of categorical logic, enabling us to establish an existential quantifier for quantum logic, and conclude that there cannot be a universal quantifier.

## 1 Introduction

Quantum logic is the study of closed subspaces of a Hilbert space [BV]. Intriguingly, this ‘logic’ is not distributive, but only satisfies the weaker axiom of orthomodularity. One of the shortcomings that has kept it from wide adoption is the lack of quantifiers. In fact, it has been called a ‘non-logic’ [Abr].

On the other hand, categorical logic [LS] can be seen as a unified framework for any kind of logic that deserves the name. It is concerned with interpreting (syntactical) logical formulae in categories with enough structure to accommodate this. An important part of it is the study of subobjects of a given object in the category at hand. Perhaps its most gratifying feature is that it gives a canonical prescription of what quantifiers should be.

The aim of this paper is to show that quantum logic is just an incarnation of categorical logic in categories like that of Hilbert spaces. In particular, we will establish an existential quantifier, and conclude that there cannot be a universal quantifier.

Section 2 first abstracts the properties of the category of Hilbert spaces that we need. This results in an axiomatisation of *(pre-)Hilbert categories* greatly resembling that of monoidal Abelian categories. In fact, any (pre-)Hilbert category embeds into the category of (pre-)Hilbert spaces itself [Heu]. Next, Section 3 starts the investigation of subobjects in Hilbert categories. It turns out that the natural objects of study are not the subobjects, but the *closed subobjects* or  $\dagger$ -*subobjects*. Section 4 then derives a functor that behaves as an existential quantifier according to categorical logic. Section 5 studies the emergent concept of orthogonality in Hilbert categories. First, it proves that  $\dagger$ -subobjects form orthomodular lattices. Second, it exhibits a tight connection between adjoint

morphisms in the base category and adjoint functors between the lattices of subobjects, the latter being important in connection to quantifiers.

### Related work

The present article should not be confused with the ‘categorical quantum logic’ of [Dun]. That work develops a type theory. Of course this is related: “every logic is a logic over a type theory” [Jac]. This paper develops the logic over ‘the type theory of Hilbert spaces’.

This paper also differs from [Har], in that the aim is explicitly a categorical logic. Another difference is that that paper restricts to those projections that have an orthocomplement, whereas we derive orthomodularity from prior assumptions (namely  $\dagger$ -kernels).

## 2 Pre-Hilbert categories

This section introduces the categories in which our study takes place, somewhat concisely. For more information we refer to [Heu].

A functor  $\dagger : \mathbf{H}^{\text{op}} \rightarrow \mathbf{H}$  with  $X^\dagger = X$  on objects and  $f^{\dagger\dagger} = f$  on morphisms is called a  $\dagger$ -*functor*; the pair  $(\mathbf{H}, \dagger)$  is then called a  $\dagger$ -*category*. Such categories are automatically isomorphic to their opposite, and the  $\dagger$ -functor witnesses this self-duality. We can consider coherence of the  $\dagger$ -functor with all sorts of structures. A morphism  $m$  in such a category that satisfies  $m^\dagger m = \text{id}$  is called a  $\dagger$ -*mono* and denoted  $\triangleright \longrightarrow$ . Likewise,  $e$  is a  $\dagger$ -*epi*, denoted  $\longrightarrow \triangleright$ , when  $ee^\dagger = \text{id}$ . A morphism is called a  $\dagger$ -*iso* when it is both  $\dagger$ -epi and  $\dagger$ -mono. Similarly, a biproduct on such a category is called a  $\dagger$ -*biproduct* when  $\pi^\dagger = \kappa$ , where  $\pi$  is a projection and  $\kappa$  an injection. This is equivalent to demanding  $(f \oplus g)^\dagger = f^\dagger \oplus g^\dagger$ . Also, an equaliser is called a  $\dagger$ -*equaliser* when it is a  $\dagger$ -mono, and a kernel is called a  $\dagger$ -*kernel* when it is a  $\dagger$ -mono. Finally, a  $\dagger$ -category  $\mathbf{H}$  is called  $\dagger$ -*monoidal* when it is equipped with monoidal structure  $(\otimes, C)$  that cooperates with the  $\dagger$ -functor, in the sense that  $(f \otimes g)^\dagger = f^\dagger \otimes g^\dagger$ , and the coherence isomorphisms are  $\dagger$ -isomorphisms.

**Definition 1** *A category is called a pre-Hilbert category when*

- *it has a  $\dagger$ -functor;*
- *it has finite  $\dagger$ -biproducts;*
- *it has (finite)  $\dagger$ -equalisers;*
- *every  $\dagger$ -mono is a  $\dagger$ -kernel; and*
- *it is symmetric  $\dagger$ -monoidal.*

Notice that a Hilbert category is self-dual (by the  $\dagger$ -functor), and therefore that it automatically has all finite colimits, too.

The category **preHilb** itself is a pre-Hilbert category whose monoidal unit is a simple generator, and so are its full subcategories **Hilb**, and **fdHilb** of finite-dimensional Hilbert spaces. Also, if **C** is a small category and **H** a pre-Hilbert category, then  $[\mathbf{C}, \mathbf{H}]$  is again a pre-Hilbert category. Working in pre-Hilbert categories can be thought of as ‘natural’ or ‘baseless’ (pre-)Hilbert space theory.

### 3 Subobjects

This section characterises closed subobjects categorically. But let us start with some easy properties of  $\dagger$ -mono’s.

**Lemma 1** *In any  $\dagger$ -category:*


- (a) *A  $\dagger$ -mono which is epi is a  $\dagger$ -iso.*
- (b) *The composite of  $\dagger$ -epi’s is again a  $\dagger$ -epi.*
- (c) *If  $X \xrightarrow{f} Y \xrightarrow{g} Z$  are such that both  $gf$  and  $f$  are  $\dagger$ -epi, so is  $g$ .*
- (d) *If  $m$  and  $n$  are  $\dagger$ -monos, and  $f$  is an iso with  $nf = m$ , then  $f$  is a  $\dagger$ -iso.*

**Proof** For (a), notice that  $ff^\dagger = \text{id}$  implies  $ff^\dagger f = f$ , from which  $f^\dagger f = \text{id}$  follows from the assumption that  $f$  is epi. For (b):  $gf(gf)^\dagger = gff^\dagger g^\dagger = g^\dagger g = \text{id}$ . And for (c):  $gg^\dagger = gff^\dagger g = gf(gf)^\dagger = \text{id}$ . Finally, consider (d). If  $f$  is iso, in particular it is epi. If both  $nf$  and  $n$  are  $\dagger$ -mono, then so is  $f$ , by (c). Hence by (a),  $f$  is  $\dagger$ -iso.  $\square$

From now on, we work in an arbitrary pre-Hilbert category **H**.

**Lemma 2** *A morphism  $m$  is mono iff  $\ker(m) = 0$ . Consequently, if  $mf = 0$  implies  $f = 0$  for all  $f$ , then  $m$  is mono.*

**Proof** Suppose  $\ker(m) = 0$ . Let  $u, v$  satisfy  $mu = mv$ . Put  $q$  to be the  $\dagger$ -coequaliser of  $u$  and  $v$ . Since  $q$  is a  $\dagger$ -epi,  $q = \text{coker}(w)$  for some  $w$ . As  $mu = mv$ ,  $m$  factors through  $q$  as  $m = nq$ . Then  $mw = nqw = n0 = 0$ , so  $w$  factors through  $\ker(m)$  as  $w = \ker(m) \circ p$  for some  $p$ . But since  $\ker(m) = 0$ ,  $w = 0$ . So  $q$  is a  $\dagger$ -iso, and in particular mono. Hence, from  $qu = qv$  follows  $u = v$ . Thus  $m$  is mono.



Conversely, if  $m$  is mono, it follows from  $m \circ \ker(m) = 0 = m \circ 0$  that  $\ker(m) = 0$ .


If  $f = 0$  whenever  $mf = 0$ , then  $\ker(m) = 0$ , so that  $m$  is mono.  $\square$

### 3.1 Factorisation

This subsection proves that any morphism  $f : X \rightarrow Y$  in a pre-Hilbert category can be factorised as an epi  $e : X \rightarrow I$  followed by a  $\dagger$ -mono  $m : I \rightarrow Y$ . (In **Hilb**, this is very easily proved concretely:  $e$  is simply the restriction of  $f$  to  $I$ , the closure of its range, and  $m$  is the isometric inclusion of  $I$  into  $Y$ .) Recall that since a pre-Hilbert category has  $\dagger$ -kernels, it automatically also has  $\dagger$ -cokernels by  $\text{coker}(f) = \ker(f^\dagger)^\dagger$ .

**Lemma 3** *Any pre-Hilbert category has a factorisation system consisting of mono's and  $\dagger$ -epi's. The factorisation is unique up to a unique  $\dagger$ -iso. Consequently, every  $\dagger$ -epi is a  $\dagger$ -cokernel of its  $\dagger$ -kernel.*

**Proof** Let a morphism  $f$  be given. Put  $k = \ker(f)$  and  $e = \text{coker}(k)$ . Since  $fk = 0$  (as  $k = \ker(f)$ ),  $f$  factors through  $e (= \text{coker}(k))$  as  $f = me$ .



We have to show that  $m$  is mono. Let  $g$  be such that  $mg = 0$ . By Lemma 2 it suffices to show that  $g = 0$ . Since  $mg = 0$ ,  $m$  factors through  $q = \text{coker}(g)$  as  $m = rq$ . Now  $qe$  is a  $\dagger$ -epi, being the composite of two  $\dagger$ -epi's. So  $qe = \text{coker}(h)$  for some  $h$ . Since  $fh = rqeh = r0 = 0$ ,  $h$  factors through  $k (= \ker(f))$  as  $h = kl$ . Finally  $eh = ekl = 0l = 0$ , so  $e$  factors through  $qe = \text{coker}(h)$  as  $q = sqe$ . But since  $e$  is a  $(\dagger)$ -epi, this means  $sq = \text{id}$ , whence  $q$  is mono. It follows from  $qg = 0$  that  $g = 0$ , and the factorisation is established.

Since  $\dagger$ -epi's are regular epi's, and hence strong epi's, functoriality of the factorisation follows from [Bor, 4.4.5]. By Lemma 1d, the factorisation is unique up to a  $\dagger$ -iso.

Finally, suppose that  $f$  is a  $\dagger$ -epi. Then both the above  $f = m \circ e$  and  $f = f \circ \text{id}$  are mono- $\dagger$ -epi factorisations of  $f$ . Hence  $f = e$  up to the unique mediating  $\dagger$ -iso  $m$ , showing that  $f = \text{coker}(\ker(f))$ .  $\square$

We just showed that any pre-Hilbert category has a factorisation system consisting of mono's and  $\dagger$ -epi's. Equivalently, it has a factorisation system of epi's and  $\dagger$ -mono's. Indeed, if we can factor  $f^\dagger$  as an  $\dagger$ -epi followed by a mono, then taking the daggers of those, we find that  $f^{\dagger\dagger} = f$  factors as an epi followed by a  $\dagger$ -mono. The combination of both factorisations yields that every morphism can be written as a  $\dagger$ -epi, followed by a monic epimorphism, followed by a  $\dagger$ -mono; this can be thought of generalising *polar decomposition*.


### 3.2 Closed subobjects, pullbacks

A *subobject* of an object  $X$  in a  $\dagger$ -category is an equivalence class of mono's  $m : M \rightarrow X$ , where  $m$  is equivalent to  $n : N \rightarrow X$  if there is an isomorphism  $f : M \rightarrow N$  satisfying  $nf = m$ . The class of subobjects of  $X$  is denoted  $\text{Sub}(X)$ . It is partially ordered by  $M \leq N$  iff there is a morphism  $f : M \rightarrow N$  with  $nf = m$ . It also has a largest element, represented by  $\text{id}_X : X \rightarrow X$ . Because a pre-Hilbert category has pullbacks,  $\text{Sub}(X)$  is in fact a meet-semilattice<sup>1</sup>, the meet of  $M$  and  $N$  being represented by the pullback of  $m$  and  $n$ . Moreover, for each  $f : X \rightarrow Y$ , pullback along  $f$  induces a meet-preserving map  $f^{-1} : \text{Sub}(Y) \rightarrow \text{Sub}(X)$ . Thus we have a functor  $\text{Sub} : \mathbf{H}^{\text{op}} \rightarrow \mathbf{MeetSLat}$ , the *inverse image functor*.

A  $\dagger$ -*subobject* is a subobject that can be represented by a  $\dagger$ -mono. We write  $\text{ClSub}(X)$  for the class of  $\dagger$ -subobjects of  $X$ . It inherits the partial ordering of  $\text{Sub}(X)$ . It can be characterised precisely when a subobject  $m$  is a  $\dagger$ -subobject, namely when there is an isomorphism  $\varphi$  such that  $m^\dagger m = \varphi^\dagger \varphi$  [Sel, 5.6].

**Lemma 4**  $\dagger$ -subobjects are stable under pullbacks.

**Proof** Let  $n : N \rightarrow Y$  and  $f : X \rightarrow Y$ . Consider their pullback  $(P, p, q)$ . Factorise  $p$  as  $p = \ker(\text{coker}(p)) \circ e$  with  $e$  epi.



Now

$$\begin{aligned} \text{coker}(n) \circ f \circ \ker(\text{coker}(p)) \circ e &= \text{coker}(n) \circ f \circ p \\ &= \text{coker}(n) \circ n \circ q \\ &= 0 \circ q = 0. \end{aligned}$$

Since  $e$  is epi,  $\text{coker}(n) \circ f \circ \ker(\text{coker}(p)) = 0$ . So there is a  $g : I \rightarrow N$  satisfying  $n \circ g = f \circ \ker(\text{coker}(p))$ . Since  $P$  is a pullback, there is a  $h : I \rightarrow P$  such that  $\ker(\text{coker}(p)) = p \circ h$ . Now  $\ker(\text{coker}(p)) = p \circ h = \ker(\text{coker}(p)) \circ e \circ h$  and  $\ker(\text{coker}(p))$  is mono, so  $e \circ h = \text{id}_I$ . Also  $p = \ker(\text{coker}(p)) \circ e = p \circ h \circ e$  and  $p$  is mono, so  $h \circ e = \text{id}_I$ . Hence  $e$  is iso, and  $p = f^{-1}(n) \in \text{ClSub}(X)$ .  $\square$

<sup>1</sup>We disregard size issues here. A  $\dagger$ -category is called  $\dagger$ -well-powered if  $\text{ClSub}(X)$  is a set for all objects  $X$  in it. Since  $\text{ClSub}(X)$  for  $X \in \mathbf{Hilb}$  is the set of closed subspaces of  $X$ ,  $\mathbf{Hilb}$  is  $\dagger$ -well-powered.

Hence every morphism  $f : X \rightarrow Y$  induces a meet-preserving map  $f^{-1} : \text{ClSub}(Y) \rightarrow \text{ClSub}(X)$ . Thus we have a functor


$$\text{ClSub} : \mathbf{H}^{\text{op}} \rightarrow \mathbf{MeetSLat},$$

that we also call the *inverse image functor* with abuse of terminology.

Recall that a *closure operation* [Bir] consists in giving for every  $m \in \text{Sub}(X)$  a  $\overline{m} \in \text{Sub}(X)$ , satisfying (i)  $m \leq \overline{m}$ , (ii)  $m \leq n \Rightarrow \overline{m} \leq \overline{n}$ , and (iii)  $\overline{\overline{m}} = \overline{m}$ .

**Lemma 5**  $m \mapsto \ker(\text{coker}(m))$  is a closure operation.

**Proof** For (i):  $\text{coker}(m) \circ m = 0$ , so  $m \leq \ker(\text{coker}(m))$ . For (ii): if  $m \leq n$ , then  $\text{coker}(m) \circ \ker(\text{coker}(m)) = 0$ ,



so  $\ker(\text{coker}(m)) \leq \ker(\text{coker}(n))$ . For (iii): since  $\ker(\text{coker}(m)) \in \text{ClSub}(X)$ , we have  $\ker(\text{coker}(\ker(\text{coker}(m)))) = \ker(\text{coker}(m))$  by Lemma 3.  $\square$

**Lemma 6** There is a reflection  $\text{Sub}(X) \xrightleftharpoons[\perp]{\ker(\text{coker}(-))} \text{ClSub}(X)$ .

**Proof** We have to prove that  $\ker(\text{coker}(m)) \leq n$  iff  $m \leq n$  for a mono  $m$  and a  $\dagger$ -mono  $n$ . By (i) of Lemma 5 we have  $m \leq \ker(\text{coker}(m))$ , proving one direction. The converse direction is just (ii) of Lemma 5.  $\square$

The previous lemma could be interpreted as a moral justification for studying the (replete) semilattice of closed subobjects instead of that of subobjects.

### 3.3 Projections

Instead of closed subobjects, it turns out we can also consider projections. A *projection* on  $X$  is a morphism  $p : X \rightarrow X$  satisfying  $p \circ p = p = p^\dagger$ . We define  $\text{Proj}(X)$  as the set of all projections on  $X$ . It is partially ordered by defining  $p \sqsubseteq q$  iff  $p \circ q = p$ .

**Proposition 1** There is an order isomorphism  $\text{ClSub}(X) \cong \text{Proj}(X)$ .

**Proof** Any closed subobject  $m$  yields a projection  $mm^\dagger$ . Conversely, any projection  $p$  gives a closed subobject  $\Im(p)$ .

Let us verify that these maps are each others inverses. Starting with a closed subobject represented by  $m$ , we end up with  $\Im(mm^\dagger)$ . Since  $m$  is  $\dagger$ -mono and

$m^\dagger$  is  $\dagger$ -epi, this is already a factorisation in the sense of Lemma 3, and hence  $\mathfrak{S}(mm^\dagger) = m$  as closed subobjects. Conversely, a projection  $p$  maps to  $ii^\dagger$ , where  $p$  factors as  $p = ie$  for an epi  $e : X \rightarrow I$  and  $\dagger$ -mono  $i = \mathfrak{S}(p)$ . By functoriality of the factorisation it follows from  $pp = p$  that  $pi = i$ . Now

$$i = pi = p^\dagger i = (ie)^\dagger i = e^\dagger i^\dagger i = e^\dagger,$$

so indeed  $ii^\dagger = ie = p$ .

Finally let us consider the order. If  $m \leq n$  as subobjects, say  $m = n\varphi$  for a  $\dagger$ -mono  $\varphi$ , then  $mm^\dagger nn^\dagger = n\varphi\varphi^\dagger n^\dagger nn^\dagger = nn^\dagger nn^\dagger = nn^\dagger$ , so indeed  $mm^\dagger \sqsubseteq nn^\dagger$ . Conversely, if  $p \sqsubseteq q$ , then  $pq = p$ , whence  $\mathfrak{S}(pq) = \mathfrak{S}(p)$ , so that indeed  $\mathfrak{S}(p) \leq \mathfrak{S}(q)$  by functoriality of the factorisation.  $\square$

Consequently, every result we derive about the partial order of closed subobjects holds for the projections and vice versa.

## 4 Existential quantifier

This section establishes an existential quantifier, *i.e.* a left adjoint to the inverse image functor that satisfies the Beck-Chevalley condition.

**Proposition 2**  $\text{ClSub}(X)$  is a lattice.

**Proof** Since we already know that  $\text{ClSub}(X)$  is a meet-semilattice, it suffices to show that it has joins and a least element. Joins follow from *e.g.* [Bor, 4.2.6].

Explicitly,  $M \vee N \xrightarrow{m \vee n} X$  is given by  $\mathfrak{S}(s)$ , where  $s = [m, n] : M \oplus N \rightarrow X$ . The closed smallest subobject, the bottom element of  $\text{ClSub}(X)$ , is given by  $0 \xrightarrow{0} X$ .  $\square$

The  $\dagger$ -mono  $m : M \rightarrowtail Y$  arising in the factorisation of a morphism  $f : X \rightarrow Y$  of  $\mathbf{H}$  is called the *(direct) image* of  $f$ , denoted  $\text{Im}(f)$ . Notice that  $\text{Im}(f)$  defines a unique  $\dagger$ -subobject, although the representing  $\dagger$ -mono is only unique up to a  $\dagger$ -iso. This  $\dagger$ -subobject is denoted  $\exists_f$ . More generally, for  $n : N \rightarrowtail X$  in  $\text{ClSub}(X)$ , we define

$$\exists_f(N) = \text{Im}(fn),$$

which gives a well-defined map  $\exists_f : \text{ClSub}(X) \rightarrow \text{ClSub}(Y)$  for any morphism  $f : X \rightarrow Y$  of  $\mathbf{H}$ .

**Theorem 1** *Let  $f : X \rightarrow Y$  be a morphism of  $\mathbf{H}$ . The map  $\exists_f : \text{ClSub}(X) \rightarrow \text{ClSub}(Y)$  is monotone and left-adjoint to  $f^{-1} : \text{ClSub}(Y) \rightarrow \text{ClSub}(X)$ . If  $g : Y \rightarrow Z$  is another morphism then  $\exists_g \circ \exists_f = \exists_{g \circ f} : \text{ClSub}(X) \rightarrow \text{ClSub}(Z)$ . Also  $\exists_{\text{id}} = \text{id}$ .*

**Proof** We follow the proof of [But, Lemma 2.5]. For monotonicity of  $\exists_f$  let  $M \leq N$  in  $\text{ClSub}(X)$ . First factorise  $n$  and then  $M \rightarrow \exists_f N$  to get the following diagram.

$$\begin{array}{ccc}
 & X & \xrightarrow{f} Y \\
 & \uparrow n & \uparrow \\
 & N & \xrightarrow{\quad} \exists_f N \\
 & \uparrow & \uparrow \\
 \curvearrowleft m & M & \xrightarrow{\quad} I
 \end{array}$$

Now  $M \twoheadrightarrow I \triangleright Y$  is an epi- $\dashv$ -mono factorisation of  $f m$ , so  $I$  represents  $\exists_f M$ , and  $\exists_f M \leq \exists_f N$ .

To show the adjunction, let  $M \in \text{ClSub}(X)$  and  $N \in \text{ClSub}(Y)$ , and consider the solid arrows in the following diagram.

$$\begin{array}{ccc}
 & X & \xrightarrow{f} Y \\
 & \uparrow & \uparrow n \\
 & f^{-1}N & \xrightarrow{\quad} N \\
 m \nearrow & & \nearrow \\
 M & \xrightarrow{\quad} \exists_f M
 \end{array}$$

If  $\exists_f M \leq N$  then the right dashed map  $\exists_f M \rightarrow N$  exists and the outer square commutes. Hence, since  $f^{-1}N$  is a pullback, the left dashed map  $M \rightarrow f^{-1}N$  exists, and  $M \leq f^{-1}N$ . Conversely, if  $M \leq f^{-1}N$ , factorise the map  $M \rightarrow N$  to get the image of  $M$  under  $f$ . In particular, this image then factors through  $N$ , whence  $\exists_f M \leq N$ .

Finally, the identity  $\exists_g \circ \exists_f = \exists_{g \circ f}$  just states how left adjoints compose.  $\square$

#### 4.1 The Beck-Chevalley condition

Recall the *Beck-Chevalley condition*: if the left square below is a pullback, then the right one must commute.

$$\begin{array}{ccc}
 \begin{array}{ccc} P & \xrightarrow{q} & Y \\ p \downarrow & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Z \end{array} & \Rightarrow & \begin{array}{ccc} \text{ClSub}(P) & \xleftarrow{q^{-1}} & \text{ClSub}(Y) \\ \exists_p \downarrow & & \downarrow \exists_g \\ \text{ClSub}(X) & \xleftarrow{f^{-1}} & \text{ClSub}(Z) \end{array} \quad (\text{BC})
 \end{array}$$

It ensures that the semantics of the existential quantifier is sound with respect to substitution. To show that our  $\exists_f$  satisfies (BC), we will assume that the monoidal unit  $C$  of our pre-Hilbert category  $\mathbf{H}$  is a simple generator. Recall



that an object  $C$  is called a *generator* when  $fx = gx$  for all  $x : C \rightarrow X$  implies  $f = g : X \rightarrow Y$ . It is called *simple* when  $\text{Sub}(C) = \{0, C\}$ . In this case, [Heu, Theorem 4.6] shows that  $\mathbf{H}$  is enriched over Abelian groups, so that we can talk of adding and subtracting morphisms.

**Lemma 7** *In a pre-Hilbert category whose monoidal unit is a simple generator, epi's are stable under pullback.*

**Proof** The proof of [Bor, Proposition 1.7.6] works verbatim.  $\square$

The previous lemma entails that  $\mathbf{H}$  is a *regular category*, and hence that all results of [But] apply. Thus, in such a category  $\mathbf{H}$  one can soundly interpret regular logic, in particular the existential quantifier.

**Theorem 2** *In a pre-Hilbert category whose monoidal unit is a simple generator, (BC) holds.*

**Proof** The proof of [But, Lemma 2.9] works verbatim.  $\square$

Also the *Frobenius identity* holds. Let  $f : X \rightarrow Y$  be a morphism of  $\mathbf{Hilb}$ . Let  $M \in \text{ClSub}(X)$  and  $N \in \text{ClSub}(Y)$ . Then  $\exists_f(M \wedge f^{-1}N) = \exists_f M \wedge N$  as  $\dagger$ -subobjects of  $Y$ . For a proof, we refer to [But, Lemma 2.6].

## 5 Orthogonality

We will now recover the orthogonal subspace construction from the  $\dagger$ -functor in any pre-Hilbert category. The idea is to mimick the fact that  $\ker(f)^\perp = \Im(f^\dagger)$  in  $\mathbf{Hilb}$ .

**Proposition 3** *There is an involutive functor  $(-)^\perp : \text{ClSub}(X)^{\text{op}} \rightarrow \text{ClSub}(X)$  determined by  $m^\perp = \ker(m^\dagger)$  for  $m \in \text{ClSub}(X)$ .*

**Proof** To show that the above definition extends functorially, let  $m, n \in \text{ClSub}(X)$  be such that  $m \leq n$ . Say that  $m$  factors through  $n$  by  $m = ni$  for  $i : M \rightarrow N$ . Then

$$m^\dagger \circ \ker(n^\dagger) = i^\dagger \circ n^\dagger \circ \ker(n^\dagger) = i^\dagger \circ 0 = 0.$$

Hence  $\ker(n^\dagger)$  factors through  $\ker(m^\dagger)$ , that is,  $n^\perp \leq m^\perp$ .

We finish the proof by showing that  $\perp$  is involutive:

$$m^{\perp\perp} = (\ker(m^\dagger))^\perp = \ker(\ker(m^\dagger)^\dagger) = \ker(\text{coker}(m)) = m.$$

Here, the last equation follows from Lemma 3.  $\square$

The functor  $(-)^\perp$  cooperates with  $\wedge$  and  $\vee$  as expected.

**Lemma 8**  *$\text{ClSub}(X)$  is an orthocomplemented lattice, that is,  $m \wedge m^\perp = 0$  and  $m \vee m^\perp = 1$  for all  $m \in \text{ClSub}(X)$ . (A fortiori, the cotuple  $[m, m^\perp]$  is a  $\dagger$ -iso.)*

**Proof** Recall that  $m \wedge m^\perp$  is defined as the  $\dagger$ -pullback

$$\begin{array}{ccc} M \wedge M^\perp & \xrightarrow{p} & M^\perp \\ \downarrow q & & \downarrow \ker(m^\dagger) \\ M & \xrightarrow{m} & X \end{array}$$

Because  $m$  is a  $\dagger$ -mono, we have  $q = m^\dagger \circ m \circ q = m^\dagger \circ \ker(m^\dagger) \circ p = 0 \circ p = 0$ . Hence  $m \wedge m^\perp = m \circ q = m \circ 0 = 0$ .

To prove the second claim, let  $f$  satisfy  $f \circ [m, m^\perp] = 0$ . Then  $f \circ m = 0$ , so  $f$  factors through  $\text{coker}(m)$  as  $f = g \circ \text{coker}(m)$ . Also  $f \circ m^\perp = 0$ , so

$$g = g \circ \ker(m^\dagger)^\dagger \circ \ker(m^\dagger) = f \circ \ker(m^\dagger) = 0,$$

whence  $f = 0$ . So, by Lemma 2,  $[m, m^\perp]$  is epi. Hence  $[m, m^\perp]$  factors as  $\text{id} \circ [m, m^\perp]$ , but also as  $(m \vee m^\perp) \circ p$ . So  $m \vee m^\perp$  must be a  $\dagger$ -iso. That is,  $m \vee m^\perp = 1$ .

Let us prove that  $[m, m^\perp]$  is also a  $\dagger$ -mono, and hence even a  $\dagger$ -iso:

$$\begin{aligned} [m, m^\perp]^\dagger \circ [m, m^\perp] &= \langle m^\dagger, \ker(m^\dagger)^\dagger \rangle \circ [m, \ker(m^\dagger)] \\ &= \begin{pmatrix} m^\dagger \circ m & m^\dagger \circ \ker(m^\dagger) \\ \ker(m^\dagger)^\dagger \circ m & \ker(m^\dagger)^\dagger \circ \ker(m^\dagger) \end{pmatrix} \\ &= \text{id}_{M \oplus M^\perp}. \end{aligned} \quad \square$$

However,  $(-)^\perp$  has poor ‘substitution properties’, as it does not commute with pullbacks. For a counterexample in **Hilb**, let  $X = \mathbb{C}^2$ ,  $Y = \mathbb{C}$ ,  $f = \pi : X \rightarrow Y : (x, y) \mapsto x$  and  $m = 0 : 0 \rightarrow Y$ . Then  $f^{-1}(m^\perp) = \mathbb{C}^2$ , but  $(f^{-1}(m))^\perp = \{(x, 0) \mid x \in \mathbb{C}\}$ .


In spite of this, a special case of “ $(-)^{\perp}$  is stable under pullbacks” still holds: we now recover orthomodularity of  $\text{ClSub}(X)$  using the previous lemma.

**Theorem 3**  $\text{ClSub}(X)$  is an orthomodular lattice, that is,  $m \vee (m^\perp \wedge n) = n$  whenever  $m \leq n \in \text{ClSub}(X)$ .


**Proof** Say that  $m$  factors through  $n$  as  $m = nd$ . Let  $(p, q)$  be the pullback of  $\ker(m^\dagger)$  along  $n$ . Then  $n \circ [q, d] = [nq, nd] = [p \circ \ker(m^\dagger), m]$ . Hence, if we can show that  $[q, d]$  is epi, this would be the factorisation of  $[p \circ \ker(m^\dagger), m]$ , and  $m \vee (m^\perp \wedge n) = n$ .

First we show that  $M \oplus (M^\perp \wedge N)$  is a pullback of  $n$  and  $[m, m^\perp]$ . Let  $f$  and  $g$  satisfy  $n \circ g = [m, \ker(m^\dagger)] \circ f$ . Then there are unique  $h_1, h_2$  making the

following diagram commute.



This makes the following into a pullback diagram.



Since isomorphisms are stable under pullback,  $[d, q]$  is iso. In particular, it is epi, and the theorem is established.  $\square$

**Corollary 1** *There cannot be right adjoints  $f^{-1} \dashv \forall_f$  for all morphisms  $f$  of  $\mathbf{H}$ , that satisfy the Beck-Chevalley condition.*

**Proof** If there would be, then  $\wedge$  would have a right adjoint in every  $\text{ClSub}(X)$  [AB, 3.4.16]. That is, there would be an implication. But the prime example **Hilb** shows that  $\text{ClSub}(X)$  is in general not a Heyting algebra.  $\square$

**Lemma 9** *The functor  $\perp: \text{ClSub}(X)^{\text{op}} \rightarrow \text{ClSub}(X)$  is an equivalence of categories. In particular, it is both left and right adjoint to its opposite  $\perp^{\text{op}}: \text{ClSub}(X) \rightarrow \text{ClSub}(X)^{\text{op}}$ .*

**Proof** This means precisely that  $m^\perp \leq n$  iff  $n^\perp \leq m$ , which holds since  $\perp$  is involutive.  $\square$

The following theorem, inspired by [Pal], provides a connection between adjoint morphisms in a pre-Hilbert category and adjoint functors between lattices of  $\dagger$ -subobjects. It explicates the relationship between  $\exists_f$  and  $\exists_{f^\dagger}$ .

**Theorem 4** *For a morphism  $f: X \rightarrow Y$ , define*

$$f^\perp = \perp_Y \circ \exists_f^{\text{op}}: \text{ClSub}(X)^{\text{op}} \rightarrow \text{ClSub}(Y).$$

*Then*

$$(f^\perp)^{\text{op}} \dashv (f^\dagger)^\perp.$$

**Proof** In general, for  $g : Y \rightarrow X$ , the adjunction  $(f^\perp)^{\text{op}} \dashv g^\perp$  means that for  $M \in \text{ClSub}(X)$  and  $N \in \text{ClSub}(Y)$ ,

$$\text{ClSub}(Y)^{\text{op}}(\perp_Y^{\text{op}} \circ \exists_f(M), N) \cong \text{ClSub}(X)(M, \perp_X \circ \exists_{f^\dagger}^{\text{op}}(N)).$$

That is,  $n \leq \ker(\exists(fm)^\dagger)$  iff  $m \leq \ker(\exists(gn)^\dagger)$ . That means that in

$$\begin{array}{ccccc} L & \xrightarrow{0} & J & & \\ \uparrow \scriptstyle q \dashv & \searrow \scriptstyle \ker(l^\dagger) & \nearrow \scriptstyle l^\dagger & \searrow \scriptstyle j^\dagger & \\ M & \xrightarrow{m} & X & \xrightarrow{g^\dagger} & Y & \xrightarrow{n^\dagger} & N \\ & \searrow \scriptstyle i & \nearrow \scriptstyle k & \searrow \scriptstyle \text{coker}(k) & & \uparrow \scriptstyle p \\ & & I & \xrightarrow{0} & K \end{array} \quad (1)$$

we must show that there is a  $p$  making the lower diagram commute iff there is a  $q$  making the upper one commute, for the special case  $g = f^\dagger$ . So, let such a  $q$  be given. Then

$$n^\dagger \circ k \circ i = n^\dagger \circ f \circ m = j \circ l^\dagger \circ m = j \circ l^\dagger \circ \ker(l^\dagger) \circ q = j \circ 0 \circ q = 0 = 0 \circ i,$$

and since  $i$  is epi,  $n^\dagger k = 0$ . Hence  $n^\dagger$  factors through  $\text{coker}(k)$  via some  $p$ . Conversely, given  $p$ , we have

$$j^\dagger \circ l^\dagger \circ m = n^\dagger \circ f \circ m = n^\dagger \circ k \circ i = p \circ \text{coker}(k) \circ k \circ i = p \circ 0 \circ i = 0 = j^\dagger \circ 0,$$

so since  $j^\dagger$  is mono,  $l^\dagger m = 0$ . Hence  $m$  factors through  $\ker(l^\dagger)$  via some  $q$ .  $\square$

In a diagram, the adjunction of the previous theorem is the following.

$$\begin{array}{ccc} \text{ClSub}(X) & \xrightarrow{\exists_f} & \text{ClSub}(Y) \\ \uparrow \scriptstyle \perp_X & \times & \downarrow \scriptstyle \perp_Y^{\text{op}} \\ \text{ClSub}(X)^{\text{op}} & \xleftarrow{\exists_{f^\dagger}^{\text{op}}} & \text{ClSub}(Y)^{\text{op}} \end{array}$$

A converse to this theorem needs some preparation, and the assumption that the monoidal unit is a simple generator.

**Lemma 10** *Let  $C$  be a simple object in a pre-Hilbert category. If  $f, g : X \rightarrow C$  satisfy  $\ker(f) \leq \ker(g)$ , then  $g = sf$  for some  $s : C \rightarrow C$ . Unless  $f = 0$ , this  $s$  is unique.*

**Proof** Consider  $\exists_f X \in \text{ClSub}(C)$ . Either  $\exists_f X = 0$ , or  $\exists_f X$  is an iso and hence a  $\dagger$ -iso since it is a  $\dagger$ -mono.

If  $\exists_f X = 0$ , then  $f = 0$ . So  $\ker(f)$  is a  $\dagger$ -iso, and since  $\ker(f) \leq \ker(g)$ , also  $\ker(g)$  is  $\dagger$ -iso, whence  $g = 0$ . Thus  $g = 0f$ .

If  $\exists_f X$  is a  $\dagger$ -iso, in particular it is epi, and so is  $f$ . It can be factorised as a  $\dagger$ -epi  $f'$  followed by a mono  $s_f$ .

$$\begin{array}{ccc} & f' \nearrow & I \searrow s_f \\ X & \xrightarrow{\quad f \quad} & C \end{array}$$

Now either  $s_f = 0$  or  $s_f$  is iso. If  $s_f = 0$  then  $\exists_f X = 0$  and hence  $f = 0$ , so that we are done by  $g = 0f$ . Hence we may assume  $s_f$  iso.

Since  $\ker(f') \leq \ker(f) \leq \ker(g)$  we are thus left with the following situation.

$$\begin{array}{ccccc} L & \xrightarrow{\ker(g)} & & f' \nearrow & C \\ \uparrow p & & & & \\ & & X & & \\ \downarrow & \xleftarrow{\ker(f')} & & g \searrow & C \\ K & & & & \end{array}$$

Now  $f' = \text{coker}(\ker(f'))$ , and

$$g \circ \ker(f') = g \circ \ker(g) \circ p = 0 \circ p = 0.$$

Hence there is a unique  $s'$  such that  $g = s' \circ f'$ . Finally, putting  $s = s' s_f^{-1}$  satisfies  $g = s' f' = s' s_f^{-1} f = s f$ .  $\square$

In a monoidal category, morphisms  $s : C \rightarrow C$  play the role of scalars, and multiplication with them is natural. As mentioned before, if  $C$  is a simple generator, then the scalars comprise an involutive field [Heu, Theorem 4.6]. The following lemma summarises some well-known (and easily proved) results.

**Lemma 11** *Let  $\mathbf{H}$  be a monoidal category. Then  $\mathbf{H}(C, C)$  is an involutive semiring that acts on  $\mathbf{H}$  by scalar multiplication as follows: for  $s : C \rightarrow C$  and  $f : X \rightarrow Y$ ,  $s \bullet f$  is defined by*

$$\begin{array}{ccc} X & \xrightarrow{s \bullet f} & Y \\ \cong \downarrow & & \uparrow \cong \\ C \otimes X & \xrightarrow{s \otimes f} & C \otimes Y \end{array}$$

*Moreover, scalar multiplication is natural, that is,  $(s \bullet g) \circ f = g \circ (s \bullet f)$ . Finally,  $s \bullet f = s \circ f$  for  $s : C \rightarrow C$  and  $f : X \rightarrow C$ .  $\square$*

Now we can state and prove a converse to Theorem 4.

**Theorem 5** *In a pre-Hilbert category whose monoidal unit is a simple generator, if  $(f^\perp)^{\text{op}} \dashv g^\perp$ , then  $g = s \bullet f^\dagger$  for a scalar  $s$ . Unless  $f = 0$ , this  $s$  is unique.*

**Proof** The adjunction of the hypothesis means that there is a  $q$  making the upper diagram in (1) commute iff there is a  $p$  making the lower one commute. So, if  $n^\dagger f m = 0$ , then  $n^\dagger k i = 0$ , and because  $i$  is epi hence  $n^\dagger k = 0$ . So  $p$  exists, whence  $q$  exists, so that  $n^\dagger g^\dagger m = j^\dagger 0 q = 0$ . Taking  $m = \ker(n^\dagger f)$  thus gives that  $\ker(n^\dagger f) \leq \ker(n^\dagger g^\dagger)$  for all  $n$ . Applying Lemma 10 yields that for all  $n : C \rightarrow Y$ , there exists  $s_n : C \rightarrow C$  such that  $n^\dagger g^\dagger = s_n n^\dagger f$ . Using Lemma 11 and dualising, this becomes: for all  $n : C \rightarrow Y$ , there is  $s_n : C \rightarrow C$  with  $g n = (s_n^\dagger \bullet f^\dagger) n$ . We will show that all  $s_n$  are in fact equal to each other (or zero). If all  $y : C \rightarrow Y$  would have  $y = 0$ , then  $Y \cong 0$ , in which case  $g = 0 \bullet f^\dagger$ . Otherwise, pick an  $y : C \rightarrow Y$  with  $y \neq 0$ . There is an  $s : C \rightarrow C$  with  $g y = (s^\dagger \bullet f^\dagger) y$ . Put  $n' = y y^\dagger n : C \rightarrow Y$  and  $n'' = \ker(y^\dagger) \circ \ker(y^\dagger)^\dagger \circ n : C \rightarrow Y$ . Then

$$\begin{aligned} n' + n'' &= [\text{id}, \text{id}] \circ ((y \circ y^\dagger \circ n) \oplus (\ker(y^\dagger) \circ \ker(y^\dagger)^\dagger \circ n)) \circ \langle \text{id}, \text{id} \rangle \\ &= [y, y^\perp] \circ [y, y^\perp]^\dagger \circ n \\ &= n. \end{aligned}$$

Moreover,

$$(s_{n'}^\dagger \bullet f^\dagger) n' = g n' = g y y^\dagger n = (s^\dagger \bullet f^\dagger) y y^\dagger n = (s^\dagger \bullet f^\dagger) n',$$

so  $s_{n'} = s$ . Finally

$$\begin{aligned} (s_{n'}^\dagger \bullet f^\dagger) n' + (s_{n''}^\dagger \bullet f^\dagger) n'' &= g n' + g n'' \\ &= g n \\ &= (s_n^\dagger \bullet f^\dagger) n \\ &= (s_n^\dagger \bullet f^\dagger) n' + (s_n^\dagger \bullet f^\dagger) n''. \end{aligned}$$

Hence  $s_n = s_{n'} = s$  for all  $n : C \rightarrow Y$ , and we have  $g n = (s^\dagger \bullet f^\dagger) n$ . But since  $C$  is a generator,  $g = s^\dagger \bullet f^\dagger$ . Reviewing our choice of  $s$  in the above proof, we see that it is unique unless  $f = 0$ .  $\square$

As a consequence, we find that, modulo scalars, the passage from morphisms  $f$  to functors  $\perp \circ \exists_f^{\text{op}}$  is one-to-one.

## A Fibred account

We can summarise our results in terms of fibred category theory [Jac]. There are fibrations  $\text{Sub}(\mathbf{H}) \rightarrow \mathbf{H}$  and  $\text{ClSub}(\mathbf{H}) \rightarrow \mathbf{H}$ . The latter is in fact a fibration of meet-semilattices by Lemma 4. The reflection of Lemma 6 is a fibred reflection. Our functor  $\exists$  of Theorem 1 is a fibred coproduct, and hence truly provides a existential quantifier.

The assignments  $\mathbf{H} \rightarrow \text{Sub}(\mathbf{H})$ ,  $X \mapsto \text{id}_X$  assemble into a fibred terminal object  $1 : \mathbf{H} \rightarrow \text{Sub}(\mathbf{H})$ , also for  $\mathbf{H} \rightarrow \text{ClSub}(\mathbf{H})$ . The fibrations  $\text{Sub}(\mathbf{H}) \rightarrow \mathbf{H}$  and  $\text{ClSub}(\mathbf{H}) \rightarrow \mathbf{H}$  admit comprehension. This means that  $1 : \mathbf{H} \rightarrow \text{Sub}(\mathbf{H})$

has a right adjoint, usually denoted by  $\{-\} : \text{Sub}(\mathbf{H}) \rightarrow \mathbf{H}$ . Indeed, if we take  $\{m : M \rightarrow X\} = M$ , then  $\text{Sub}(\mathbf{H})(\text{id}_X, m) \cong \mathbf{H}(X, \{m\})$ .

In fact, the fibration  $\text{ClSub}(\mathbf{H}) \rightarrow \mathbf{H}$  is a bifibration by Theorem 1 and [Jac, 9.1.2] – notice that the Beck-Chevalley condition is not needed for this. Thus,  $\text{ClSub}(\mathbf{H})^{\text{op}} \rightarrow \mathbf{H}^{\text{op}}$ ,  $(m : M \twoheadrightarrow X) \mapsto X$  is also a fibration. The following proposition shows that orthogonality can be extended to a functor between fibrations, but it is not a fibred functor, basically because it does not commute with pullback.

**Proposition 4**  $(-)^{\perp}$  extends to a functor  $\text{ClSub}(\mathbf{H})^{\text{op}} \rightarrow \text{ClSub}(\mathbf{H})$  satisfying

$$\begin{array}{ccc} \text{ClSub}(\mathbf{H})^{\text{op}} & \xrightarrow{(-)^{\perp}} & \text{ClSub}(\mathbf{H}) \\ \downarrow & & \downarrow \\ \mathbf{H}^{\text{op}} & \xrightarrow{(-)^{\dagger}} & \mathbf{H} \end{array} \quad (2)$$

However, it is not a fibred functor.

**Proof** We can understand  $(-)^{\perp}$  as a functor  $\text{ClSub}(\mathbf{H})^{\text{op}} \rightarrow \text{ClSub}(\mathbf{H})$  by extending its action on morphisms as follows. Let  $(f, g)$  be a morphism  $m \rightarrow n$ , that is, let  $f : X \rightarrow Y$  and  $g : M \rightarrow N$  satisfy  $fm = ng$ . We are to define a morphism  $(f, g)^{\perp} : n^{\perp} \rightarrow m^{\perp}$ , that is, a pair  $f^{\perp} : Y \rightarrow X$  and  $g^{\perp} : N^{\perp} \rightarrow M^{\perp}$  satisfying  $f^{\perp} \circ n^{\perp} = m^{\perp} \circ g^{\perp}$ . Put  $f^{\perp} = f^{\dagger}$ . Then

$$m^{\dagger} \circ f^{\dagger} \circ n^{\perp} = g^{\dagger} \circ n^{\dagger} \circ \ker(n^{\dagger}) = g^{\dagger} \circ 0 = 0,$$

so there is a  $g^{\perp}$  such that  $f^{\dagger} \circ n^{\perp} = \ker(m^{\dagger}) \circ g^{\perp} = m^{\perp} \circ g^{\perp}$ . It must be

$$g^{\perp} = (m^{\perp})^{\dagger} \circ m^{\perp} \circ g^{\perp} = \ker(m^{\dagger})^{\dagger} \circ f^{\dagger} \circ n^{\perp} = \text{coker}(m) \circ f^{\dagger} \circ \ker(n^{\dagger}).$$

This explicitly defines the functor  $(-)^{\perp} : \text{ClSub}(\mathbf{H})^{\text{op}} \rightarrow \text{ClSub}(\mathbf{H})$ . It makes the square (2) commute. Now, a morphism  $(f, g) : m \rightarrow n$  of  $\text{ClSub}(\mathbf{H})$  is Cartesian (over  $f$ ) iff  $f = ngm^{\dagger} = nn^{\dagger}fmm^{\dagger}$ . Consequently, the morphism  $(f, g)^{\perp} : n^{\perp} \rightarrow m^{\perp}$  in  $\text{ClSub}(\mathbf{H})^{\text{op}}$  is Cartesian iff

$$f^{\dagger} = \ker(m^{\dagger}) \circ \ker(m^{\dagger})^{\dagger} \circ f^{\dagger} \circ \ker(n^{\dagger}) \circ \ker(n^{\dagger})^{\dagger}.$$

Thus,  $(-)^{\perp}$  is a fibred functor iff  $f^{\dagger} = \ker(m^{\dagger})\ker(m^{\dagger})^{\dagger}f^{\dagger}\ker(n^{\dagger})\ker(n^{\dagger})^{\dagger}$  whenever  $f = nn^{\dagger}fmm^{\dagger}$  for any morphism  $f$  and  $\dagger$ -mono's  $m$  and  $n$ .

Finally, we come to our counterexample. Take  $m = \kappa_M$  for  $M \neq 0$ ,  $f = mm^{\dagger}$  and  $n = \text{id}_{M \oplus M}$ . Then  $f = mm^{\dagger} = mm^{\dagger}mm^{\dagger} = nn^{\dagger}fmm^{\dagger}$ . But  $\ker(m^{\dagger}) = \kappa'$  so  $\ker(m^{\dagger})^{\dagger} = \pi'$  and  $\ker(n^{\dagger}) = 0$ , so

$$\ker(m^{\dagger}) \circ \ker(m^{\dagger})^{\dagger} \circ f^{\dagger} \circ \ker(n^{\dagger}) \circ \ker(n^{\dagger})^{\dagger} = \kappa' \circ \pi' \circ f^{\dagger} \circ 0 = 0 \neq f^{\dagger}.$$

Hence  $(-)^{\perp}$  is not a fibred functor.  $\square$

## References

- [AB] Steve Awodey and Andrej Bauer. Introduction to categorical logic.
- [Abr] Samson Abramsky. Temperley-lieb algebra: From knot theory to logic and computation via quantum mechanics. In Goong Chen, Louis Kauffman, and Sam Lomonaco, editors, *Mathematics of Quantum Computing and Technology*, pages 415–458. Taylor and Francis, 2007.
- [Bir] Garrett Birkhoff. *Lattice Theory*. American Mathematical Society, 1948.
- [Bor] Francis Borceux. *Handbook of Categorical Algebra 1: Basic Category Theory*. Encyclopedia of Mathematics and its Applications 50. Cambridge University Press, 1994.
- [But] Carsten Butz. Regular categories and regular logic. BRICS Lecture Series LS-98-2, 1998.
- [BV] Garrett Birkhoff and John Von Neumann. The logic of quantum mechanics. *Annals of Mathematics*, 37:823–843, 1936.
- [Dun] Ross Duncan. *Types for Quantum Computing*. PhD thesis, Oxford University Computer Laboratory, 2006.
- [Har] John Harding. Orthomodularity in dagger biproduct categories. *submitted to International Journal of Theoretical Physics*, 2008.
- [Heu] Chris Heunen. An embedding theorem for Hilbert categories. *submitted to Theory and Applications of Categories*, 2008.
- [Jac] B. Jacobs. *Categorical Logic and Type Theory*. Number 141 in Studies in Logic and the Foundations of Mathematics. North Holland, 1999.
- [LS] Joachim Lambek and Phil Scott. *Introduction to higher order categorical logic*. Cambridge University Press, 1986.
- [Pal] Paul H. Palmquist. Adjoint functors induced by adjoint linear transformations. *Proceedings of the American Mathematical Society*, 44(2):251–254, 1974.
- [Sel] Peter Selinger. Idempotents in dagger categories. In *QPL 2006*, pages 87–101, 2006.